

## Original Investigation

# Sleep-Deprived Young Drivers and the Risk for Crash

## The DRIVE Prospective Cohort Study

Alexandra L. C. Martiniuk, MSc, PhD; Teresa Senserrick, PhD; Serigne Lo, PhD; Ann Williamson, PhD; Wei Du, PhD; Ronald R. Grunstein, MD, PhD; Mark Woodward, PhD; Nick Glozier, MBBS, PhD; Mark Stevenson, PhD, MPH; Robyn Norton, PhD; Rebecca Q. Ivers, MPH, PhD

← Editorial page 606

**IMPORTANCE** Short sleep duration is common in adolescents and young adults, and short sleep duration is a risk factor for motor vehicle crash.

**OBJECTIVE** To assess the association between hours of sleep and the risk for motor vehicle crash, including the time of day of crash and types of crash (single, multiple vehicle, run off road, and intersection).

**DESIGN** Prospective cohort study.

**SETTING** New South Wales, Australia.

**PARTICIPANTS** Questionnaire responses were obtained from 20 822 newly licensed drivers aged 17 to 24 years. Participants held a first-stage provisional license between June 2003 and December 2004 prospectively linked to licensing and police-reported crash data, with an average of 2 years of follow-up. Analyses were conducted on a subsample of 19 327 participants for which there was full information.

**EXPOSURE** Sleeping 6 or fewer hours per night.

**MAIN OUTCOMES AND MEASURES** The main outcome variable was police-reported crash. Multivariable Poisson regression models were used to investigate the role of sleep duration on the risk for crash.

**RESULTS** On average, those who reported sleeping 6 or fewer hours per night had an increased risk for crash compared with those who reported sleeping more than 6 hours (relative risk [RR], 1.21; 95% CI, 1.04-1.41). Less weekend sleep was significantly associated with an increased risk for run-off-road crashes (RR, 1.55; 95% CI, 1.21-2.00). Crashes for individuals who had less sleep per night (on average and on weekends) were significantly more likely to occur between 8 PM and 6 AM (RR, 1.86; 95% CI, 1.11-3.13, for midnight to 5:59 AM and RR, 1.66; 95% CI, 1.15-2.39, for 8:00 PM to 11:59 PM).

**CONCLUSIONS AND RELEVANCE** Less sleep per night significantly increased the risk for crash for young drivers. Less sleep on weekend nights increased the risk for run-off-road crashes and crashes occurring in the late-night hours. This provides rationale for governments and health care providers to address sleep-related crashes among young drivers.

**Author Affiliations:** The George Institute for Global Health, University of Sydney, Sydney, Australia (Martiniuk, Lo, Woodward, Norton, Ivers); Woolcock Institute, University of Sydney, Sydney, Australia (Grunstein); Brain and Mind Research Institute, University of Sydney, Sydney, Australia (Glozier); Sydney Medical School, University of Sydney, Sydney, Australia (Lo); Transport and Road Safety, University of New South Wales, Sydney, Australia (Senserrick, Williamson); Neuroscience Research Australia, Sydney, Australia (Du); Monash University Accident Research Center, Melbourne, Australia (Stevenson); University of Toronto, Toronto, Ontario, Canada (Martiniuk).

**Corresponding Author:** Alexandra Martiniuk, MSc, PhD, The George Institute for Global Health, University of Sydney, PO Box M201, Missenden Road, Sydney, NSW 2050, Australia (amartiniuk@george.org.au).

*JAMA Pediatr.* 2013;167(7):647-655. doi:10.1001/jamapediatrics.2013.1429  
Published online May 20, 2013.

Every day, more than 3000 people die in motor vehicle crashes around the world, amounting to more than 1.3 million motor vehicle-related deaths worldwide each year—and between 20 million and 50 million more are injured and often left disabled for life.<sup>1</sup> Motor vehicle crashes also have a huge economic impact worldwide, responsible for up to 3% of a country's gross national product costs per year.<sup>1</sup> In the United States, it is estimated that drowsy driving is responsible for 20% of all motor vehicle crashes, meaning that in the United States alone, drowsy driving may cause 1 million crashes, 50 000 injuries, and 8000 deaths each year.<sup>2</sup> Young drivers are involved in a disproportionate number of these crashes.<sup>3</sup> Adolescents and young adults are at particular risk for sleep deprivation. There are many lifestyle factors that lead to reduced sleep hours in adolescents and young adults, including biological development, academic responsibilities, socialization, and employment, as well as increased use of electronic communication devices.<sup>4-6</sup> Population-based data indicate young adults sleep less than older adults.<sup>7</sup> Moreover, identical periods of experimental sleep deprivation result in greater sleepiness and impaired attention in younger adults compared with healthy elderly people.<sup>8</sup> This higher biological sleep need reflects developmental changes in sleep, arousal, and circadian regulation.<sup>4,9</sup>

Worldwide, adolescents frequently say they get inadequate sleep and experience daytime sleepiness,<sup>10</sup> including 60% of US youth aged 17 to 24 years.<sup>11</sup> Complaints of sleepiness are associated with lapses in attention, slowed reaction time, impairments in judgment, difficulty regulating emotions including increased aggression, and risky behavior; these effects are magnified when alcohol is also involved.<sup>4,5,9,10,12-22</sup> Young drivers (aged <25 years) are at the highest risk for sleep-related motor vehicle crashes.<sup>4,13,23-25</sup> Australian data show that although only 20% of their driving occurs at night,<sup>24</sup> 60% of young driver deaths happen at night,<sup>26</sup> with a US study showing 37% of such deaths occur on weekend nights.<sup>25</sup> While all drivers have a greater risk for crashing at night, existing data suggest young drivers are at even greater risk, potentially with lack of sleep a contributing factor. For drivers of all ages, estimates in the United States, United Kingdom, and Australia report that between 5% and 30% of crashes are attributed to fatigue.<sup>3,24,27-31</sup> Young drivers report driving while tired, with 51% of US young drivers reporting driving while drowsy and 9% reporting nodding off while driving in the past year.<sup>32</sup> Not only are they more likely to have sleep-related crashes, these crashes are more likely to be fatal compared with other crash causes.<sup>33</sup> In one study, young driver crashes classified as fatigue related tended to be single-vehicle crashes if they occurred during the night, but they were more often head-on crashes during the day.<sup>24</sup>

Importantly, to our knowledge, previous studies of crash risk due to sleepiness have not focused on young drivers and have mainly had descriptive,<sup>34</sup> case-control, or retrospective cohort designs.<sup>27,35,36</sup> While these studies have provided valuable information, retrospective designs risk biased exposure information. For example, information on sleep collected after a crash could be affected by social desirability response bias. Also, data on fatigue-related crashes are often subjectively reported by police at the crash scene. Another important limitation of many previous studies is the inability to control for important confounders

such as driving exposure, psychological disturbances, urban/rural status, alcohol use, and risk taking. A prospective study allows the opportunity to collect information on chronic sleepiness and other possible confounders, such as alcohol use and mental health, before the crash occurs. Therefore, we used a large prospective cohort study to determine the role of sleep on the risk for future motor vehicle crashes. Specifically, we aimed to determine crash risk in young drivers according to the number of sleep hours per night, the relationship between weekday and weekend sleep hours and crash risk, and relationships for different types and times of crashes among those who crashed.

## Methods

### The Drive Study

Data were derived from the DRIVE Study, a prospective cohort of 20 822 drivers aged 17 to 24 years in New South Wales (NSW), Australia. Detailed information on participant heterogeneity and study methods have been reported elsewhere.<sup>37</sup> In brief, all drivers resident in NSW aged 17 to 24 years holding a first-stage provisional license between June 2003 and December 2004 were invited to participate. Follow-up data were collected until the a priori selected completion date of December 31, 2005. An online baseline survey queried sleep hours (weekends and weekdays), as well as demographics, driving exposure, driver training, drug and alcohol use, risky driving, and sensation seeking. All respondents gave consent for their survey data to be linked prospectively to data held by the state jurisdictional authority, the Roads and Traffic Authority, including information about police-reported crashes. Average follow-up time was 2 years. Ethics committees of the University of Sydney and NSW Health approved the study.

### Study Sample

Of 131 000 eligible new drivers, 20 822 (16%) participated in the DRIVE Study. Analyses were conducted on a subsample of 19 327 participants for which there was full information on sleep duration (1087 participants had missing data), risky driving behaviors (318 missing), and average weekly driving hours (90 missing). In this sample, there were 1383 crashes (of which 280 were single vehicle and 305 run off road).

### Study Outcome: Crash

The outcome variable was police-recorded crash as a driver. In NSW, a crash is recorded by police when a person is killed or injured; when the crash is a hit and run; when damage to property is more than A\$500; when 1 or more of the drivers was reported to be driving under the influence of alcohol; or if a vehicle involved in the crash was towed away. We classified this primary outcome as binary (ie, none vs 1 or more crashes), as well as the secondary outcomes of crash times and crash types (eg, a type of crash vs all other crashes). Thus, the secondary analyses had a reduced sample size, including only those who crashed, to determine the unique role sleep duration may have played.

### Study Exposure: Sleep

Total sleep hours were assessed at the time of the baseline survey by asking: "During the past month, about how long have you

slept for each night (weekday hours) and (weekend hours)”—a standard assessment used in large epidemiological studies reporting on the consequences of sleep duration (eg, National Health and Nutrition Examination Survey).<sup>38</sup> Values provided were rounded to the nearest integer value. The weighted average of weekday and weekend sleep hours was used to determine average sleep duration. Sleep was categorized according to 3 classifications: average sleep hours per week, average weekday sleep hours, and average weekend (Saturday and Sunday) sleep hours. We contrasted sleep hours within each classification as 6 or fewer hours vs more than 6 hours. Previous research has focused on a 5-hour cutoff<sup>33,27</sup> when exploring crash risk, but a 6-hour cutoff for other health risks.<sup>11,14,15</sup> Because sleeping 6 or fewer hours is expected to affect more of the population, this cutoff was chosen owing to its anticipated public health impact.<sup>11,14,15</sup>

### Confounders

Confounding variables were collected at the time of the baseline survey, including age group (17, 18-19, or 20-24 years), sex, average weekly driving hours ( $\leq 2$ , 3-4, 5-9, or  $\geq 10$  hours), remoteness of residence (urban, inner region, or rural),<sup>39</sup> drinking behavior (Alcohol Use Disorders Identification Test score  $< 7$  or  $\geq 7$ ),<sup>40</sup> and risky driving behaviors (adapted from Donovan and Jessor<sup>41</sup> and the University of Otago Young Driver Pilot Study<sup>42</sup>), such as speeding (categorized as low, medium, and high frequency), self-harm (yes/no),<sup>43,44</sup> crash history (yes/no), drug-use habit (never,  $\leq 1$ /month, 2-4 times/month, 2-3 times/week, and  $\geq 4$  times a week),<sup>45</sup> sensation seeking (low, medium, and high),<sup>46</sup> and Kessler Psychological Distress Scale score (low, moderate, high, and very high).<sup>47,48</sup> The information collected on confounders refers to the 1 month preceding the baseline survey, except for the question about self-harm, which refers to the 12 months preceding the baseline survey.

### Statistical Analyses

Relative risks (RRs) for the association between sleep duration (hours) and crash involvement were calculated using Poisson regression. Poisson regression models were used, which allows for an offset that takes into account time in the study since subject entry and exit from the study, and thus time at risk for crash varied. All models controlled for prior crashes. To examine the association between sleep hours and crashes, we forced all confounders into the model, while for sleep hours and crash types we forced only age group, sex, average weekly driving hours, remoteness of residence, and drinking behavior owing to smaller sample sizes. Interaction terms involving sleep hours were added individually to regression models, and a likelihood ratio test was used to compare model with interaction terms vs model without interaction terms. We intended to retain these interactions for *P* values less than .05; however, no interaction was significant. SAS version 9.1 was used for all analyses.<sup>49</sup>

## Results

Young drivers' sleep hours by crash risk factors are presented in the **Table** for weekday and weekend nights. Older drivers (aged 18-19 and 20-24 years) were more likely to sleep 6 or fewer

hours per night than younger drivers (aged 17 years); 10% of 17-year-old subjects, 15% of 18- to 19-year-old subjects, and 17% of 20- to 24-year-old subjects slept 6 or fewer hours a night. There was no difference in sleep hours by remoteness of residence. Men and women reported sleeping approximately an equal number of hours, except women slept less than men on weekends. Drivers who slept fewer hours per night drove more hours than those who slept more. Risky driving, sensation seeking, self-harm, higher Kessler Psychological Distress Scale score, and greater drug and alcohol intake were reported more often by individuals who obtained less sleep.

**Figure 1** presents the RRs for crash by average hours of sleep per day. In unadjusted models, shorter sleep duration was significantly associated with increased crash risk for 6 or fewer hours vs more than 6 hours of sleep a night (RR, 1.36; 95% CI, 1.17-1.57). Adjusted models demonstrated a slightly attenuated risk for crash (RR, 1.21; 95% CI, 1.04-1.41). Shorter sleep duration per night was significantly related to multiple vehicle crashes ( $\leq 6$  hours vs  $> 6$  hours of sleep a night; RR, 1.30; 95% CI, 1.11-1.53). Run-off-road-type crashes neared significance in the relationship with shorter sleep duration ( $\leq 6$  hours vs  $> 6$  hours of sleep a night; RR, 1.37; 95% CI, 1.00-1.89). Less sleep per night ( $\leq 6$  hours vs  $> 6$  hours) was significantly associated with crashes occurring in 2 periods: midnight to 5:59 AM (RR, 1.86; 95% CI, 1.11-3.13) and 8:00 PM to 11:59 PM (RR, 1.66; 95% CI, 1.15-2.39).

**Figure 2** presents the RRs for crash by average hours of sleep per weekday night. In unadjusted models, shorter sleep duration was significantly associated with increased crash risk (RR, 1.20; 95% CI, 1.05-1.38). Adjusted models demonstrated a slightly attenuated risk for crash (RR, 1.12; 95% CI, 0.97-1.29). Shorter sleep duration per night was significantly related to multiple-vehicle crashes ( $\leq 6$  hours vs  $> 6$  hours of sleep a night; RR, 1.19; 95% CI, 1.02-1.38). Less sleep per night ( $\leq 6$  hours vs  $> 6$  hours) was not significantly associated with crashes occurring in any particular period.

**Figure 3** presents the RRs for crash by average hours of sleep per weekend night. In unadjusted models, shorter sleep duration was significantly associated with increased crash risk (RR, 1.24; 95% CI, 1.10-1.39). Adjusted models demonstrated risk for crash, but they were nonsignificant for 6 or fewer hours vs more than 6 hours of sleep per night (RR, 1.08; 95% CI, 0.95-1.23). Shorter sleep duration per weekend night was significantly related with run-off-road-type crashes ( $\leq 6$  hours vs  $> 6$  hours of sleep a night; RR, 1.55; 95% CI, 1.21-2.00). Less sleep per night ( $\leq 6$  hours vs  $> 6$  hours) was significantly associated with crashes occurring in 2 periods: midnight to 5:59 AM (RR, 1.78; 95% CI, 1.13-2.80) and 8:00 PM to 11:59 PM (RR, 1.42; 95% CI, 1.03-1.94). Results for sleeping 5 or fewer hours a night were similar to the findings for those sleeping 6 or fewer hours per night.

## Discussion


This study demonstrated that sleeping 6 or fewer hours per night, similar to 5 or fewer hours, is associated with an increased risk for motor vehicle crashes among young drivers. This increased risk remained even after controlling for confounders including exposure to driving.<sup>12-18</sup> Reduced average sleep hours per night

Table. Description of Factors by Average Hours of Sleep Per Night<sup>a</sup>

	Average Sleep Hours/Night, No. (%)				Total (N = 19 327)
	Weekdays		Weekends		
	≤6 (n = 2759)	>6 (n = 16 568)	≤6 (n = 3963)	>6 (n = 15 364)	
<b>Age group, y</b>					
17	1083 (39.3)	8336 (50.3)	1785 (45.0)	7634 (49.7)	9419 (48.7)
18-19	1138 (41.3)	6057 (36.6)	1667 (42.1)	5528 (36.0)	7195 (37.2)
20-24	538 (19.5)	2175 (13.1)	511 (12.9)	2202 (14.3)	2713 (14.0)
<b>Sex</b>					
Male	1300 (47.1)	7525 (45.4)	1596 (40.3)	7229 (47.1)	8825 (45.7)
Female	1459 (52.9)	9043 (54.6)	2367 (59.7)	8135 (52.9)	10 502 (54.3)
<b>Average driving, h</b>					
<2	519 (18.8)	3306 (20.0)	484 (12.2)	3341 (21.8)	3825 (19.8)
3-4	423 (15.3)	3187 (19.2)	583 (14.7)	3027 (19.7)	3610 (18.7)
5-9	719 (26.1)	5006 (30.2)	1241 (31.3)	4484 (29.2)	5725 (29.6)
≥10	1098 (39.8)	5069 (30.6)	1655 (41.8)	4512 (29.4)	6167 (31.9)
<b>Risk-taking level</b>					
Low	789 (28.6)	5921 (35.7)	825 (20.8)	5885 (38.3)	6710 (34.7)
Medium	900 (32.6)	5422 (32.7)	1300 (32.8)	5022 (32.7)	6322 (32.7)
High	1070 (38.8)	5225 (31.5)	1838 (46.4)	4457 (29.0)	6295 (32.6)
<b>Residential remoteness</b>					
Urban	2234 (81.0)	12 161 (73.4)	2886 (72.8)	11 509 (74.9)	14 395 (74.5)
Inner region	444 (16.1)	3629 (21.9)	898 (22.7)	3175 (20.7)	4073 (21.2)
Rural	81 (2.9)	778 (4.7)	179 (4.5)	680 (4.4)	859 (4.4)
<b>Self-harm</b>					
Yes	351 (12.7)	1148 (6.9)	469 (11.8)	1030 (6.7)	1499 (7.8)
No	2329 (84.4)	15 187 (91.7)	3430 (86.6)	14 086 (91.7)	17 516 (90.6)
Unknown	11 (0.4)	233 (1.4)	64 (1.6)	248 (1.6)	312 (1.6)
<b>Crash history</b>					
Yes	909 (33.0)	4804 (29.0)	1453 (36.7)	4260 (27.7)	5713 (29.6)
No	1839 (66.7)	11 713 (70.7)	2495 (63.0)	11 057 (72.0)	13 552 (70.1)
Unknown	11 (0.4)	51 (0.3)	15 (0.4)	47 (0.3)	62 (0.3)
<b>Kessler score</b>					
Low	486 (17.6)	5096 (30.8)	898 (22.7)	4684 (30.5)	5582 (28.9)
Moderate	924 (33.5)	6480 (39.1)	1455 (36.7)	5949 (38.7)	7404 (38.3)
High	898 (32.6)	3909 (23.6)	1153 (29.1)	3654 (23.8)	4807 (24.9)
Very high	442 (16.0)	1023 (6.2)	445 (11.2)	1020 (6.6)	1467 (7.6)
Unclassified	9 (0.3)	60 (0.4)	12 (0.3)	57 (0.4)	69 (0.4)
<b>Sensation score</b>					
Low	705 (25.6)	5400 (32.6)	814 (20.5)	5291 (34.4)	6105 (31.6)
Medium	821 (29.8)	5417 (32.7)	1110 (28.0)	5128 (33.4)	6238 (32.3)
High	1209 (43.8)	5582 (33.7)	1991 (50.2)	4800 (31.2)	6791 (35.1)
Unclassified	24 (0.9)	169 (1.0)	48 (1.2)	145 (0.9)	193 (1.0)
<b>Drinks/wk</b>					
<7	2340 (84.8)	14 422 (87.1)	2969 (74.9)	13 793 (89.8)	16 762 (86.7)
>7	419 (15.2)	2141 (12.9)	994 (25.1)	1566 (10.2)	2560 (13.3)
Unknown	0 (0)	5 (0)	0 (0)	5 (0)	5 (0)
<b>Drug-use habit, times/mo</b>					
Never	2491 (90.3)	15 452 (93.3)	3444 (86.9)	14 499 (94.4)	17 943 (92.8)
1	182 (6.6)	823 (5.0)	360 (9.1)	645 (4.2)	1005 (5.2)
2-4	52 (1.9)	172 (1.0)	110 (2.8)	114 (0.7)	224 (1.2)
2-3	17 (0.6)	32 (0.2)	26 (0.7)	23 (0.2)	49 (0.3)
≥4 times/wk or more	14 (0.5)	27 (0.2)	16 (0.4)	25 (0.2)	41 (0.2)
Unknown	3 (0.1)	62 (0.4)	7 (0.2)	58 (0.4)	65 (0.3)

<sup>a</sup> All adjusted models included age, sex, exposure to driving, remoteness of residence, and alcohol; for crashes, we added the supplementary variables of previous crash, self-harm, Kessler Psychological Distress Scale score, sensation-seeking scale, and drug use.

Figure 1. Relative Risk for Crash by Average Sleep Hours per Day (>6 Hours vs ≤ 6 Hours)




\*Adjusted analyses of all crashes by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, drinking behavior (Alcohol Use Disorders Identification Test score), risky driving behaviors, self-harm, drug use, sensation seeking, and

psychological distress. Adjusted analyses of crash types by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, and driving behavior owing to smaller sample sizes.

was also significantly associated with crashes occurring between 8 PM and 6 AM. Importantly, this study also found that reduced sleep hours on the weekends was significantly associated with an increased risk for run-off-road crashes and crashes occurring between 8 PM and 6 AM. Our findings offer a potential improvement over past research of sleep and crash that have focused on a 5-hour cutoff, which is a less common level of reduced sleep hours, and have been retrospective or case-control studies and thus more likely to have had important biases.<sup>24,27,50,51</sup> Previous studies have observed greater magnitudes of crash risk for those who are sleepy (RR, 8.2; 95% CI, 3.4-19.7), have had fewer than 5 hours of sleep (RR, 2.7; 95% CI, 1.4-5.4), drive between 2 AM and 5 AM (RR, 5.6; 95% CI, 1.4-22.7),<sup>27</sup> and those reporting they were falling asleep (RR, 14.2; 95% CI, 1.4-14.7).<sup>36</sup> These relatively large RRs from earlier studies are bounded by wide confidence intervals indicating smaller sample sizes compared with the current data. The lower RRs observed in the current study compared with previous literature may be owing to weaknesses in prior study design. Alternatively, it may be owing to the concept of drowsiness, which differs from reported hours of sleep, and may have

more to do with changes in factors such as sleep pattern or length of driving. Differences may also be owing to the inclusion of all crashes not just injury crashes in the current study. In the DRIVE Study sample, about 13% of the cohort reported sleeping 6 or fewer hours per night, consistent with previous estimates of 15%.<sup>14,52,53</sup> In the DRIVE Study, drivers who slept fewer hours per night were more likely to be older, report a high number of weekly driving hours, engage in risky driving, be high-sensation seekers, engage in self-harm, and have higher Kessler Psychological Distress Scale scores, and greater drug and alcohol use. It is possible that those who slept less drove more because of work or school commuting, although sleep hours did not differ by remoteness of residence. Previous authors have suggested that the nighttime driving restrictions within graduated driver licensing may deal indirectly with some portion of this excess risk, but it has been argued that these restrictions do not resolve morning risks when driving to school or work or afternoon hours returning home, when adolescents can be similarly fatigued.<sup>54</sup> However, our study only observed an increased risk from 8 PM to 6 AM. This highlights a need to further investigate nighttime driving

Figure 2. Relative Risk for Crash by Average Sleep Hours per Weekday (>6 Hours vs ≤6 Hours)



\*Adjusted analyses of all crashes by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, drinking behavior (Alcohol Use Disorders Identification Test score), risky driving behaviors, self-harm, drug use, sensation seeking, and

psychological distress. Adjusted analyses of crash types by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, and driving behavior owing to smaller sample sizes.


restrictions.<sup>55,56</sup> In our study data, the hypothesized increased risk for early-morning driving crashes (eg, driving to school/work, which we would expect after 6 AM) was not observed.

This study has significant strengths. First, to our knowledge, no prospective study has investigated the relationship between sleep and motor vehicle crash. Second, this study used police-recorded crashes, providing an objective outcome measure with little loss to follow-up. Third, this study enrolled a large number of participants, controlled for several potential confounders, and investigated sleep hours by weekdays and weekends, as well as more specific crash types and times. These findings may help increase awareness of the impact of reduced sleep hours on crash risk and highlight subgroups of young drivers and times of day for targeted intervention. Owing to feasibility in this large, prospective cohort study, average measures of sleep per night on weekdays and weekends and Alcohol Use Disorders Identification Test scores for alcohol use were taken at baseline; however, we recognize these are not as ideal as measures of sleep deprivation and alcohol intoxication at the time of the crash. For instance, asking about usual or average sleep may not reflect vari-

ability of sleep, and we cannot determine whether crashes occurred on days of little sleep. However, previous research indicates that crashes can be affected both by proximal sleep deprivation (eg, the day of the crash) and more distal effects of chronic sleep loss (eg, months preceding the crash), as well as the effects of sleep loss on the acquisition of a new skill such as driving.<sup>25</sup> More objective methods to measure sleep (eg, actigraphy) were unavailable for this study, having an unknown effect on the results. Also, some analyses, such as single-vehicle crash analyses, were limited by small numbers even though the study enrolled more than 20 000 young drivers; thus, we observed less precision around the magnitude of risk for these analyses. Additionally, previous research has demonstrated that single-vehicle crashes are less likely to be reported to police, which may have occurred in this study. Lastly, this study enrolled participants via an online, voluntary questionnaire and was not designed to be generalizable. However, a heterogeneous population of young drivers was represented in this study.

A range of measures may prevent crashes due to sleepiness or reduce their seriousness. Changes to road design (eg,

Figure 3. Relative Risk for Crash by Average Sleep Hours per Weekend (>6 Hours vs ≤ 6 Hours)



\*Adjusted analyses of all crashes by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, drinking behavior (Alcohol Use Disorders Identification Test score), risky driving behaviors, self-harm, drug use, sensation seeking, and psychological distress. Adjusted analyses of crash types by sleep hours controlled for time in the study, prior crashes, age group, sex, average weekly driving hours, remoteness of residence, and driving behavior owing to smaller sample sizes.

tactile road edges and divided highways), as well as education campaigns, may help reduce crash risk.<sup>57,58</sup> Using a rest stop (RR, 0.5), drinking coffee (RR, 0.5), and playing the radio while driving (RR, 0.6) have been shown to be significantly protective against crashes, at least in the short term.<sup>36</sup> In Australia, New Zealand, the United States, and several other countries, extended campaigns on the short 30-minute powernap are run by road authorities.<sup>32,59,60</sup>

Although residual confounding may remain, in particular by acute alcohol intoxication, reduced sleep hours is a risk factor for crashes independent of other covariates.<sup>61,62</sup> Young drivers sleeping a few hours per night also had other risk factors for crashes, thus interventions may be well placed to address multiple risk factors simultaneously. Future research could also evaluate the effects of harm-minimization strategies at parties

and other youth entertainment events, such as alcohol sales restrictions, and legal and peer dynamics education along with education to reduce drowsy driving.<sup>63</sup> Young drivers should be a key focus because this group experiences more impairment in alertness, mood, and physical performance compared with older age groups with similar sleep deprivation.<sup>8</sup>

Our current paper addresses several of the research needs highlighted during the recent Hopkinton Conference on Future Directions in Fatigue and Safety Research including time of day effects, fatigue, and risk taking,<sup>64</sup> as well as the use of prospective study designs.<sup>65</sup> This study provides rationale for addressing sleep deprivation among young drivers<sup>66</sup>—in particular highlighting increased risks for crash with 6 or fewer hours of sleep per night, weekend run-off-road crashes, and crashes between 8 PM and 6 AM.

ARTICLE INFORMATION

Accepted for Publication: October 31, 2012.

Published Online: May 20, 2013.

doi:10.1001/jamapediatrics.2013.1429.

**Author Contributions:** Dr Martiniuk is the guarantor for this study. All authors had full access to all the data and take full responsibility for the integrity of this study and its analyses. *Study concept and design:* Senserrick, Williamson,

Woodward, Stevenson, Norton, Ivers. *Acquisition of data:* Martiniuk, Woodward, Stevenson, Ivers. *Analysis and interpretation of data:* Lo, Williamson, Du, Grunstein, Woodward, Glozier, Ivers.

*Drafting of the manuscript:* Martiniuk, Senserrick, Williamson, Glozier, Stevenson, Ivers.

*Critical revision of the manuscript for important intellectual content:* Martiniuk, Senserrick, Lo, Williamson, Du, Grunstein, Woodward, Glozier, Norton, Ivers.

*Statistical analysis:* Martiniuk, Lo, Williamson, Du, Woodward, Glozier.

*Obtained funding:* Woodward, Stevenson, Norton, Ivers.

*Administrative, technical, and material support:* Martiniuk, Stevenson, Norton.

*Study supervision:* Woodward, Glozier, Stevenson, Ivers.

**Conflict of Interest Disclosures:** Drs Martiniuk, Senserrick, Williamson, Du, Stevenson, and Ivers received salary funding from the National Health and Medical Research Council of Australia during the writing of this article. Dr Martiniuk's work is funded by a fellowship from the University of Sydney.

**Funding/Support:** The DRIVE Study was funded by the National Health and Medical Research Council of Australia, Roads and Traffic Authority of New South Wales, NRMA Motoring and Services, NRMA-ACT Road Safety Trust, New South Wales Health, and the Motor Accidents Authority of New South Wales.

**Disclaimer:** No funding agency had any role in the design, analysis, or interpretation of the data.

**Additional Information:** Data on motor vehicle crashes were collected by the Roads and Traffic Authority of New South Wales.

## REFERENCES

- World Health Organization. Global status report on road safety 2009. [www.who.int/violence\\_injury\\_prevention/road\\_safety\\_status/2009/en/index.html](http://www.who.int/violence_injury_prevention/road_safety_status/2009/en/index.html). Accessed June 9, 2010.
- Colten HR, Altevogt BM, eds. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. Washington, DC: Board on Health Sciences Policy, National Academies Press; 2006.
- World Health Organization. Youth and road safety, April 2007. [www.who.int/violence\\_injury\\_prevention/publications/road\\_traffic/youth\\_road\\_safety/en/index.html](http://www.who.int/violence_injury_prevention/publications/road_traffic/youth_road_safety/en/index.html). Accessed June 9, 2010.
- Dahl RE. Biological, developmental, and neurobehavioral factors relevant to adolescent driving risks. *Am J Prev Med*. 2008;35(3)(suppl):S278-S284.
- Ohayan MM, Roberts RE, Zulley J, Smirene S, Priest RG. Prevalence and patterns of problematic sleep amongst older adolescents. *J Am Child Adolesc Psychiatry*. 2000;39(12):1549-1556.
- Van den Bulck J. Adolescent use of mobile phones for calling and for sending text messages after lights out: results from a prospective cohort study with a one-year follow-up. *Sleep*. 2007;30(9):1220-1223.
- Krueger PM, Friedman EM. Sleep duration in the United States: a cross-sectional population-based study. *Am J Epidemiol*. 2009;169(9):1052-1063.
- Duffy JF, Willson HJ, Wang W, Czeisler CA. Healthy older adults better tolerate sleep deprivation than young adults. *J Am Geriatr Soc*. 2009;57(7):1245-1251.
- Roberts RE, Roberts CR, Duong HT. Chronic insomnia and its negative consequences for health and functioning of adolescents: a 12-month prospective study. *J Adolesc Health*. 2008;42(3):294-302.
- Carskadon MA. Patterns of sleep and sleepiness in adolescents. *Pediatrics*. 1990;17(1):5-12.
- Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep patterns and predictors of disturbed sleep in a large population of college students. *J Adolesc Health*. 2010;46(2):124-132.
- Barrett PR, Horne JA, Reyner LA. Alcohol continues to affect sleepiness related driving impairment, when breath alcohol levels have fallen to near-zero. *Hum Psychopharmacol*. 2004;19(6):421-423.
- Pack AI, Pack AM, Rodgman E, Cucchiara A, Dinges DF, Schwab CW. Characteristics of crashes attributed to the driver having fallen asleep. *Accid Anal Prev*. 1995;27(6):769-775.
- Bartlett DJ, Marshall NS, Williams A, Grunstein RR. Sleep health New South Wales: chronic sleep restriction and daytime sleepiness. *Intern Med J*. 2008;38(1):24-31.
- Gibson ES, Powles AC, Thabane L, et al. "Sleepiness" is serious in adolescence: two surveys of 3235 Canadian students. *BMC Public Health*. 2006;6:116.
- Singleton RA Jr, Wolfson AR. Alcohol consumption, sleep, and academic performance among college students. *J Stud Alcohol Drugs*. 2009;70(3):355-363.
- Liu X, Zhao Z, Jia C, Buysse DJ. Sleep patterns and problems among Chinese adolescents. *Pediatrics*. 2008;121(6):1165-1173.
- Killgore WD, Lipizzi EL, Kamimori GH, Balkin TJ. Caffeine effects on risky decision making after 75 hours of sleep deprivation. *Aviat Space Environ Med*. 2007;78(10):957-962.
- McKenna BS, Dickinson DL, Orff HJ, Drummond SP. The effects of one night of sleep deprivation on known-risk and ambiguous-risk decisions. *J Sleep Res*. 2007;16(3):245-252.
- Killgore WD, Balkin TJ, Wesensten NJ. Impaired decision making following 49 h of sleep deprivation. *J Sleep Res*. 2006;15(1):7-13.
- Lucidi F, Russo PM, Mallia L, Devoto A, Lauriola M, Violani C. Sleep-related car crashes: risk perception and decision-making processes in young drivers. *Accid Anal Prev*. 2006;38(2):302-309.
- O'Brien EM, Mindell JA. Sleep and risk-taking behavior in adolescents. *Behav Sleep Med*. 2005;3(3):113-133.
- McConnell CF, Bretz KM, Dwyer WO. Falling asleep at the wheel: a close look at 1,269 fatal and serious injury-producing crashes. *Behav Sleep Med*. 2003;1(3):171-183.
- Dobbie K; Australian Transport Safety Bureau. Fatigue-related crashes: an analysis of fatigue-related crashes on Australian roads using an operational definition of fatigue. 2002. [www.infrastructure.gov.au/roads/safety/publications/2002/pdf/Fatigue\\_related\\_sum.pdf](http://www.infrastructure.gov.au/roads/safety/publications/2002/pdf/Fatigue_related_sum.pdf). Accessed August 22, 2010.
- Wolfson AR, Carskadon MA. Sleep schedules and daytime functioning in adolescents. *Child Dev*. 1998;69(4):875-887.
- Australian Transport Safety Bureau. *Road Safety In Australia: A Publication Commemorating World Health Day 2004*. Canberra, Australia: Australian Transport Safety Bureau; 2004.
- Connor J, Norton R, Ameratunga S, et al. Driver sleepiness and risk of serious injury to car occupants: population based case control study. *BMJ*. 2002;324(7346):1125.
- Fell D. *Safety Update: Problem Definition and Countermeasure Summary: Fatigue 1994*. RUS No. 5. New South Wales, Australia: New South Wales Road Safety Bureau.
- Fell DL, Black B. Driver fatigue in the city. *Accid Anal Prev*. 1997;29(4):463-469.
- Fell D. The road to fatigue: circumstances leading to fatigue accidents. In: Hartley L, ed. *Fatigue and Driving*. Bristol, PA: Taylor and Francis Inc; 1995.
- Horne J, Reyner L. Vehicle accidents related to sleep: a review. *Occup Environ Med*. 1999;56(5):289-294.
- National Sleep Foundation of America website. [www.sleepfoundation.org](http://www.sleepfoundation.org). Accessed June 10, 2010.
- Bunn TL, Slavova S, Struttman TW, Browning SR. Sleepiness/fatigue and distraction/inattention as factors for fatal versus nonfatal commercial motor vehicle driver injuries. *Accid Anal Prev*. 2005;37(5):862-869.
- Radun I, Summla H. Sleep-related fatal vehicle accidents: characteristics of decisions made by multidisciplinary investigation teams. *Sleep*. 2004;27(2):224-227.
- Stutts JC, Wilkins JW, Scott Osberg J, Vaughn BV. Driver risk factors for sleep-related crashes. *Accid Anal Prev*. 2003;35(3):321-331.
- Cummings P, Koepsell TD, Moffat JM, Rivara FP. Drowsiness, counter-measures to drowsiness, and the risk of a motor vehicle crash. *Inj Prev*. 2001;7(3):194-199.
- Ivers RQ, Blows SJ, Stevenson MR, et al. A cohort study of 20,822 young drivers: the DRIVE Study methods and population. *Inj Prev*. 2006;12(6):385-389.
- Gangwisch JE, Heymsfield SB, Boden-Albala B, et al. Short sleep duration as a risk factor for hypertension: analyses of the first National Health and Nutrition Examination Survey. *Hypertension*. 2006;47(5):833-839.
- Chen HY, Ivers RQ, Martiniuk AL, et al. Risk and type of crash among young drivers by rurality of residence: findings from the DRIVE Study. *Accid Anal Prev*. 2009;41(4):676-682.
- Babor TF, de la Fuente JR, Saunders J, et al. *AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for Use In Primary Health Care*. Geneva, Switzerland: World Health Organization; 1992.
- Donovan JE, Jessor R. *Young Adult Driving Questionnaire*. Boulder, CO: Colorado Institute of Behavioral Science; 1992.
- Begg D, Brookland R, Hope J, Langley J, Broughton J. New Zealand drivers study: developing a methodology for conducting a follow-up study of newly licensed drivers. *J Safety Res*. 2003;34(3):329-336.
- Patton GC, Harris R, Carlin JB, et al. Adolescent suicidal behaviours: a population-based study of risk. *Psychol Med*. 1997;27(3):715-724.

44. Martiniuk AL, Ivers RQ, Glozier N, et al. Self-harm and risk of motor vehicle crashes among young drivers: findings from the DRIVE Study. *CMAJ*. 2009;181(11):807-812.
45. Brener ND, Kann L, Kinchen SA, et al. Methodology of the youth risk behaviour surveillance system. *MMWR Recomm Rep*. 2004;53(rr-12):1-13.
46. Zuckerman M, Kuhlman DM, Joireman J, et al. A comparison of the three structural models for personality: the big three, the big five, and the alternative five. *J Pers Soc Psychol*. 1993;65(4):757-768. doi:10.1037/0022-3514.65.4.757.
47. Australian Bureau of Statistics. Information paper: use of the Kessler Psychological Distress Scale in ABS health surveys, Australia, 2001. [www.abs.gov.au/ausstats/abs@.nsf/ProductsbyReleaseDate/4D5BD324FE8B415FCA2579D500161D57](http://www.abs.gov.au/ausstats/abs@.nsf/ProductsbyReleaseDate/4D5BD324FE8B415FCA2579D500161D57). Accessed April 19, 2013.
48. Martiniuk ALC, Ivers RQ, Glozier N, et al. Does psychological distress increase the risk for motor vehicle crashes in young people? findings from the DRIVE Study. *J Adolesc Health*. 2010;47(5):488-495.
49. SAS Institute Inc. *Statistical Analysis Software (SAS) Version 9.1*. Cary, NC: SAS Institute Inc; 2002.
50. Gander PH, Marshall NS, Harris RB, Reid P. Sleep, sleepiness and motor vehicle accidents: a national survey. *Aust N Z J Public Health*. 2005;29(1):16-21.
51. Liu GF, Han S, Liang DH, et al. Driver sleepiness and risk of car crashes in Shenyang, a Chinese northeastern city: population-based case-control study. *Biomed Environ Sci*. 2003;16(3):219-226.
52. Dorofaeff TF, Denny S. Sleep and adolescence: do New Zealand teenagers get enough? *J Paediatr Child Health*. 2006;42(9):515-520.
53. Paine SJ, Gander PH, Harris RB, Reid P. Prevalence and consequences of insomnia in New Zealand: disparities between Maori and non-Maori. *Aust N Z J Public Health*. 2005;29(1):22-28.
54. Keating DP, Halpern-Felsher BL. Adolescent drivers: a developmental perspective on risk, proficiency, and safety. *Am J Prev Med*. 2008;35(3)(suppl):S272-S277.
55. McCartt AT, Teoh ER, Fields M, Braitman KA, Helling LA. Graduated licensing laws and fatal crashes of teenage drivers: a national study. *Traffic Inj Prev*. 2010;11(3):240-248.
56. Williams AF. Contribution of the components of graduated licensing to crash reductions. *J Safety Res*. 2007;38(2):177-184.
57. Haworth N. Countering driver fatigue. In: Langford J, Fildes B, eds. *Australasian Road Safety Handbook, Volume 1. Sydney, Australia: Austroads*;46-53.
58. Hatfield J, Murphy S, Job RFS, Du W. The effectiveness of audio-tactile lane-marking in reducing various types of crash: a review of evidence, template for evaluation, and preliminary findings from Australia. *Accid Anal Prev*. 2009;41(3):365-379.
59. VicRoads Australia. Fatigue and road safety 2010. [www.vicroads.vic.gov.au/Home/SafetyAndRules/SafetyIssues/Fatigue/FatigueAndRoadSafety.htm](http://www.vicroads.vic.gov.au/Home/SafetyAndRules/SafetyIssues/Fatigue/FatigueAndRoadSafety.htm). Accessed April 19, 2013.
60. New Zealand Ministry of Transport National Road Safety Committee. Driver fatigue strategy 2009. [www.transport.govt.nz/about/publications/Documents/DriverStrategicFrameworkamend3.pdf](http://www.transport.govt.nz/about/publications/Documents/DriverStrategicFrameworkamend3.pdf). Accessed June 10, 2010.
61. Banks S, Catchside P, Lack L, Grunstein RR, McEvoy RD. Low levels of alcohol impair driving simulator performance and reduce perception of crash risk in partially sleep deprived subjects. *Sleep*. 2004;27(6):1063-1067.
62. Wilson RJ, Fang M, Cooper PJ, Beirness DJ. Sleepiness among night-time drivers: relationship to blood alcohol concentration and other factors. *Traffic Inj Prev*. 2006;7(1):15-22.
63. Youth Safe. Safe celebrating 2010. [www.youthsafe.org/](http://www.youthsafe.org/). Accessed June 10, 2010.
64. Horrey WJ, Noy YI, Folkard S, Popkin SM, Howarth HD, Courtney TK. Research needs and opportunities for reducing the adverse safety consequences of fatigue. *Accid Anal Prev*. 2011;43(2):591-594.
65. Williamson A, Lombardi DA, Folkard S, Stutts J, Courtney TK, Connor JL. The link between fatigue and safety. *Accid Anal Prev*. 2011;43(2):498-515.
66. Simons-Morton BG. Parent involvement in novice teen driving: rationale, evidence of effects, and potential for enhancing graduated driver licensing effectiveness. *J Safety Res*. 2007;38(2):193-202.